

The Effect of Soil Moisture Content in Soil Microbial Biomass Loblolly Pine Stand

Shrijana Duwadi¹, Ryan Nadel¹, Yucheng Feng², Lori G. Eckhardt¹

¹ Forest Health Dynamics Laboratory, School of Forestry and Wildlife Sciences, Auburn University, Auburn, Alabama, USA; ² College of Agriculture, Auburn University, Auburn, Alabama, USA



Results




Fig 5: Bivariate fit of MB-C by SMC during Winter 2016.




Fig 5: Bivariate fit of MB-N by SMC during Winter 2016.




Fig 6: Bivariate fit of MB-C by SMC during Spring 2016.




Fig 6: Bivariate fit of MB-N by SMC during Spring 2016.




Fig 7: Bivariate fit of MB-C by SMC during Summer 2016.




Fig 7: Bivariate fit of MB-N by SMC during Summer 2016.




Fig 8: Bivariate fit of MB-C by SMC during Fall 2016.




Fig 8: Bivariate fit of MB-N by SMC during Fall 2016.

Summary

- In every season, soil moisture was observed to be lower than the previous season.
- MB was not affected by the SMC in the winter and the spring, while MB was affected by SMC in the summer and the fall.
- Soil microbes might be limited due to low moisture availability in the summer and the fall.

