

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Stinger\*

EPA Reg. No. 62719-73

### Control of Sicklepod and Other Susceptible Broadleaf Weeds in Southern Pine Seedbeds in Forest Nurseries

For distribution and use only in the states of Alabama, Arkansas, Georgia, Louisiana,  
Mississippi, North Carolina, South Carolina, Tennessee, Texas and Virginia

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Stinger before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Stinger according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Stinger.

#### Directions for Use

Stinger\* herbicide may be applied over-the-top of loblolly pine, slash pine, and longleaf pine to control sicklepod and other susceptible broadleaf weeds in southern pine seedbeds in forest nurseries. Apply as a broadcast or spot treatment from May through July when weeds are actively growing. Refer to the product label for Stinger for a complete listing of weeds controlled.

#### Application Timing

**General broadleaf weed control:** For best results, apply when weeds are small and actively growing.

**Sicklepod:** For best results, apply after the majority of basal leaves have emerged.

#### Application

Apply at a broadcast rate of 1/4 to 1/2 pt per acre in a spray volume of 20 or more gallons per acre. Application may be made any time after May 1, but some needle curling may occur if applied during active conifer growth. When making spot applications, use a calibrated boom, or if a hand-held sprayer is used, care should be taken to apply the spray uniformly and at a rate equivalent to a broadcast application (See guidelines in the label for Stinger for use of hand-held sprayers). Otherwise, do not use more than 1/5 fl oz (1 tsp.) of Stinger per gallon of spray and direct spray onto weeds and avoid spraying pine seedlings whenever possible.

#### Precautions:

- Application of Stinger during active growth of conifers may cause some needle curling.
- Do not use surfactants or crop oils in spray mixtures as the potential for tree injury in the form of needle curling may be increased.

\*Trademark of Dow AgroSciences LLC

A2A / Stinger / FPL-Supp / 05-12-99

D06-043-006

EPA-accepted: 05-03-99

Initial printing