

SPECIMEN



**For Use as a Preemergent Weed Control Herbicide in Turfgrasses, Landscape
or Grounds Maintenance, Noncropland Areas and Ornamental Production**

ACTIVE INGREDIENT

pendimethalin, N-(1-ethylpropyl)-3,4-dimethyl-2, 6-dinitrobenzenamine 38.7%

INERT INGREDIENTS: 61.3%

TOTAL 100.0%

(1 gallon contains 3.8 lbs. of microencapsulated pendimethalin in an aqueous carrier.)

EPA Reg. No. 241-416

EPA Est. No. _____

**KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See next page for First Aid instructions and additional Precautionary Statements

For more information, please visit our web site
www.turffacts.com

BASF
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709

 **BASF**
The Chemical Company

SPRAYING INSTRUCTIONS

Uniformly apply with properly calibrated ground equipment in suggested spray volumes of 20-200 gpa for ornamental applications to uniformly treat the area with a spray pressure of 25 to 50 psi. Maintain continuous agitation during spraying with good mechanical or bypass agitation. Avoid overlaps that will increase rates above those recommended. Avoid application when winds may cause drift.

Avoid unintentional contact of spray solution with driveways, stone, wood, or other porous surfaces. Rinse immediately with water to avoid staining. Avoid mechanically scrubbing until surface area is thoroughly rinsed using a heavy spray of water.

INSTRUCTIONS AND RESTRICTIONS¹ IN PRODUCTION ORNAMENTALS

Site	Application Instructions and Restrictions
Newly-Transplanted Field-Grown Nursery Stock ^{2,3}	<ol style="list-style-type: none"> DO NOT make over-the-top applications at time of field transplanting. Use shielded sprayer until plantings have been established for one (1) year or more in the field. DO NOT apply until transplants have been watered and soil has been thoroughly packed and settled around transplants. Care must be taken to ensure there are no cracks in the soil where Pendulum® AquaCap™ herbicide could come into contact with the roots. DO NOT apply during bud swell, bud break or at time of first flush of new growth. Direct sprays away from graphed or budded tissue on transplants at all times.
Newly-Transplanted Container-Grown Nursery Stock ^{2,3}	<ol style="list-style-type: none"> DO NOT apply until transplants have been watered and soil has been thoroughly packed and settled around transplants. Care must be taken to ensure there are no cracks in the soil where PENDULUM AquaCap could come into contact with the roots. For container grown ornamentals, delay first application of the product to bareroot liners for two (2) weeks after transplanting. DO NOT apply during bud swell, bud break or at time of first flush of new growth. Direct sprays away from graphed or budded tissue on transplants at all times.
Established Container, or Field-Grown Nursery Stock ^{2,3}	<ol style="list-style-type: none"> DO NOT apply during bud swell, bud break or at time of first flush of new growth. Apply as a directed or over-the-top spray. If newly budded or graphed rootstock, make an application using a shielded sprayer. Care must be taken to ensure there are no cracks in the soil where PENDULUM AquaCap could come into contact with the roots.
Bare Ground for Container Placement	<ol style="list-style-type: none"> Apply to soil then water in (including mulch, gravel, wood chips, or other permeable base), replace containerized ornamentals onto pad.
Greenhouses, shadehouses or other enclosed structures.	DO NOT APPLY in greenhouses, shadehouses or other enclosed structures.

¹ Plant only those desirable plant species listed on this label into soil treated the previous season with **Pendulum AquaCap** or injury may occur.

² It is recommended that before treating a large number of plants, spray a few plants and observe for 1-2 months for plant damage prior to full-scale application.

³ **DO NOT** treat plants grown for food or feed. **DO NOT** use treated plants for food or feed.

Refer to **Table 3. Application Rates For Weed Control In Production Ornamentals.**