### FOR USE ONLY WITHIN THE STATE OF OKLAHOMA

### RESTRICTED USE PESTICIDE

Due to toxicity to aquatic invertebrate animals.

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.



# DIMILIN® 25W

NSECT GROWTH REGULATOR EPA REG. No. 400-465

Active Ingredients: (% by weight)

## KEEP OUT OF REACH OF CHILDREN CAUTION

### THIS LABEL IS IN ADDITION TO THE STANDARD FEDERAL LABEL PRINTED ON THE CONTAINER

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **DIRECTIONS FOR USE**

#### CHRISTMAS TREES AND PINE TREE NURSERIES

**To control Nantucket Pine Tipmoth:** Apply 4 ounces of DIMILIN 25W per acre beginning in spring when first instar larvae are present or 75% of pupal cases are empty. Various methods may be used, i.e. twig sampling, pheromone traps, degree days, etc. to determine DIMILIN 25W application at peak emergence. Repeat applications as needed for subsequent generations. Do not make more than 3 applications per season.

Aerial Application: Apply the recommended amount of DIMILIN 25W in 0.5 to 2.0 gallons of water per acre for uniform coverage of foliage.

Ground Application: Apply the recommended amount of DIMILIN 25W sufficient water for uniform, full coverage of the top 1/3 of the tree (10-200 gal.). Low gallonage (5-10 gal/acre), high pressure sprays are preferred for better distribution and coverage. Do not apply spray to runoff.

Follow all applicable directions, restrictions, and precautions on the EPA registered label. This label must be in the possession of the user at the time of the pesticide application.

002