

Reference

Forest Nursery Pests

USDA Forest Service Ag Handbook 680



Nursery Insect Pests

- Leaf/foliage feeders or sap suckers
- Damage seedlings by feeding on roots, stem, shoot and buds
- Causes seedling mortality, reduces seedling grade, culls

Major Nursery Insect Pests

- Tarnish Plant Bug - *Lygus lineolaris*
 - Attacks a wide variety of economically important herbaceous plants, vegetable crops, commercial flower plants, fruit trees, and nursery stock.
 - Lygus bugs occur in all Canadian provinces, the continental United States and most of Mexico.
 - Approximately 50% of loblolly pine seedlings in one southern forest nursery was damaged by Lygus bugs (South 1986)



Tarnish plant bug – Lygus Bug

Tarnish Plant Bug

- The insects over winters as adults in dead weeds, leaf litter, under tree bark, nursery margins, ditch banks, and road rights-of-way.
- Insects become active in early spring and feed on newly developing buds and shoots. Most nursery damage occurs from mid-April to late June.
- Oviposition is restricted to composite host plants (non-conifers) where eggs are deposited at the base of the leaf blade.

Tarnish Plant Bug

- After 7-10 days, yellowish-green nymphs emerge and begin feeding. The life cycle is completed in three to four weeks.
- There are two to three generations per year.
- At least 385 host plants have been recorded for *Lygus* with most in the Rosidae and Asteridae families.
- The insect also attacks pine seedlings which are severely damaged.

Tarnish Plant Bug

- Adults and nymphs of *Lygus* feed by sucking plant juices and inject into the plant a watery saliva to aid in the breakdown of plant tissues.
- The feeding causes terminal growth to be distorted thereby reducing plant growth. Damaged by *Lygus* feeding has been called "crazy cotton", "stop-back", "bush- head", "bushy-top."
- Symptoms appear within a few weeks after feeding and apical dominance is lost and weak multiple leaders appear.

Tarnish Plant Bug

- In conifer seedlings, terminal needles are thicker and shorter and the tip is often curled
- The removal of preferred host plants from edges of nurseries and destruction of favorable overwintering sites will help to reduce the damages caused by *Lygus*.
- Weed hosts include butterweed, fleabane, goldenrod, vetch, dock, and dog fennel.
- Several insecticides are available to control populations of *Lygus*.



Insecticide treated bed

Non-treated bed



“Bushy-Top” symptoms of Lygus bug feeding damage

Major Nursery Insect Pests

- Lesser Cornstalk borer
Elasmopalpus lignosellus
- Found throughout the southern US
- Larva feed upon the seedling stem; mortality, secondary pathogens enter

Lesser cornstalk borer

- Look for wounds just below to just above the ground level.
- Bark may be completely or partially removed.
- Partially girdled seedlings may have a gall/swelling on stem.
- Seedlings turn chlorotic, orange, and die; remain standing upright or tip over.

Lesser cornstalk borer

- Larva less than an inch in length.
- Pale green with brown bands/stripes.
- May produce silk tunnels in soil.
- Wriggle furiously when handled.
- Rare to find the larva.
- Adult moths, more commonly observed.
- Moth-like in color, fly erratically above seedlings, about an inch in length.



Lesser cornstalk borer larva



Lesser cornstalk borer adult

Lesser cornstalk borer

- Insect has 2-4 generations per year.
- Late summer, all life stages are present.
- Adult moths emerge from the soil in late spring, mate and female deposits eggs at base of seedlings.
- Eggs hatch within 7 days and larva feed on lower stem, or subterranean roots.
- Larva feed 3 wks, pupate in soil, emerge, mate lay eggs.
- Over winter as both larva and pupae in soil.

Lesser cornstalk borer

- Cover crops, sandy soils and drought favor LCB activity.
- Insect prefers corn, but it also feeds on beans, cowpeas, crabgrass, Johnson grass, peas, peanuts, sorghum, soybeans, and wheat.
- Cultivation promotes, rather than retards, injury by insect. Damage is less under no-tillage cropping systems which is attributed to increased soil moisture and the presence of decaying organic matter.
- Insecticides available to use against LCB; chlorpyrifos

Regeneration Weevils

Pales weevil
(*Hylobius pales*)

Pitch eating weevil
(*Pachylobius picivorus*)





UGA0949010b

Pales Weevil - Found throughout S.E.

Biology

- Adults attracted to fresh resin odor and they invade recently cut over areas and eggs laid in pine stump roots.
- Eggs hatch in 5-10 days and the larvae feed on inner bark of dead roots - not a problem.
- Weevils pupate in chip cocoons and adults fly to new seedlings feed on tender bark of pine branches.
- Newly planted seedlings = girdle stem and kill
- 2 generations/yr but adults present year round in south.



Pales Weevil Damage



Pales Weevil Damage

Why is timing important?

Weevils attracted to cut timber areas & lay eggs at base of stumps. They reproduce in large numbers.

Immediate replanting means you are putting the seedlings (500/acre) into a "sea" of weevils.

Waiting one year allows the insects time to disperse from the area.

Management

Cultural - alter replanting times

Chemical- Insecticides (Pounce)


- Prior to lifting

- At time of packing

- In the field (spot)

Pine Tip Moth

- Most damaging to pine plantations and to wild pine seedlings in open areas.
- In the South and Southeast, the favored hosts are loblolly and shortleaf pine. Pitch, Virginia



Pine Tip Moth

- Pine tip moth injures the growing shoots of young pines.
- The larva bores into and feeds on inner tissues of the buds and shoots.
- Such feeding severs the conductive tissue and causes death of the shoot
- Shoot injury occurs primarily in the first 5 years and decreases as the tree reaches about 10 feet (3 m) in height and the crown closes.



Pine Tip Moth

- Insect overwinters as a pupa within damaged shoots of the host trees.
- On warm days, as early as February in the South, adult moths emerge, mate, and lay eggs on the current season's shoots and conelets.
- Eggs may take as long as 30 days to hatch if cool weather follows egg laying in the spring; later in the summer, eggs hatch in only 5 to 10 days.
- Newly hatched larvae feed on the surface of new growth and cause shallow injuries, or they may bore into the needle fascicles.
- Later the larvae move to the shoot tips.

Minor Nursery Insect Pests

- White Grubs

Phyllophaga spp.

- 100 different species and other genera of white grubs include: *Diploptaxis*, *Dichelonyx*, *Serica*, & *Cotalpa*

Symptoms:

- Seedling foliage turns brown, seedling roots cut off, 3-5 mm gouges in larger roots, tap root severed, smaller roots missing.
- Appears in newly established nurseries, 2-3 yr post fumigation and in outplanting areas with sod.
- Found throughout eastern United States

Identification



- Grubs always "C" shaped
- Found in soil near roots
- Roots appear sparse or have chewed upon look

Identification



James L. Castner, U. Fla. Ent. Dep.

Adults large brown or black beetles May beetles, June bugs, Green June Beetles, Japanese beetles.

Life Cycle:

- Adults strongly attracted to lights & often found in pools of water
- Nocturnal feeders on hardwoods, especially oaks and can defoliate stands of oaks

Life cycle: 3-yr, sometimes only 2-yr in southern US

- Eggs laid in summer in soil near seedlings
- Larvae feed on roots until fall, then burrow deep in soil, hibernate in the soil.
- Spring, move back up to feed on roots
- Cycle repeated two more years,
- Larvae grow bigger each year, cause increasing damage

Life Cycle

- Larvae complete growth third spring.
- Pupate in soil for a few weeks.
- Adults emerge from pupal case but remain in soil until next spring
- Fly to oaks, feed, mate, return to seedlings to lay eggs for next generation of white grubs.

Damage

- Very injurious to seedlings
- Damage worst within a few 100 yards of *Quercus* sp
- Adults do not fly far from food to lay eggs
- 1 larva/sq ft causes serious damage

Damage

- After fumigation (or seedling establishment) damage is minimal first year
- Becomes more serious 2nd year
- Very severe third year as larvae grow larger and eat more each year
- Normally beds must be treated every 3 yr

White Grub Management

Keep adult food plants such as *Quercus* sp away from Nursery. Beetles are lazy and poor fliers.

Insecticides

- Fumigate beds with MBr/Chl before sowing
- Granular insecticides sometimes effective
- Dipping of seedlings in insecticide reduces damage after outplanting - high risk sites.







White Grub Management

Fumigate every 3-4 yrs

Spot treat with Discus (Imidacloprid + Cyfluthrin)

Minor - Mole crickets



Mole crickets:

May be important locally, but not always recognized.

Southern Pest, virtually all of Florida, the southern half of Alabama. However, recently found in Anniston. Not a problem in cold climates

Mole cricket inactive until soil reaches 60 F. Feed at night, in upper 1 inch of soil

Damage - Two types

Southern Mole cricket, Tunneling disrupts roots, uproots seedlings but does not feed on roots



Tawney mole cricket

Feeds on roots, damage can be serious.
Is common in Florida, rarer as move north.



Mole cricket damage in nursery bed

Mole Cricket Biology

- Adults over-winter in deep soil burrows.
- Move up in soil profile when temperatures are right, feed.
- Adults lay eggs in soil chambers, which hatch in June.
- Pupate in soil for a few weeks.
- Two peaks of feeding activity.
- First in March/April when overwintering adults begin feeding.
- Second in Sept/Oct when the new generation of nymphs feed and the adults continue to feed.
- Second peak most damaging. There is only one generation per year

Management - Insecticides

1st peak of activity- Control optional due to low numbers. Usually when it is observed.

2nd Peak – Use Orthene, Dursban (chlorpyrifos products)

Biological control. Parasitic wasps and nematodes available

Populations monitored/controlled using artificial cricket callers

Minor insect: Cutworms

Several
species of
Noctuidae



Distributed all over U.S. with the most damaging in Lake States and south

High populations can destroy 1000's of seedlings in a few Weeks

Damage symptoms include cut off needles and seedlings clipped at soil level

Chemical sprays and fumigation are effective
chlorpyrifos, Asana, Discus



Figure 49-1—Cutworm damage on young conifer seedlings. Note clipped needles.



Figure 49-2—Dingy cutworm larva (left) and pupa.




Figure 49-3—Dingy cutworm adult.

Summary for Insecticides

- The use of Asana as the primary tool against Lygus is a good choice.
- The use of chlorpyrifos as the primary tool against LCB is ok but this is a poor choice for Lygus.
- There is probably a better choice than chlorpyrifos for LCB.
- Use the 24c for Pales only as intended.
- Discus white grubs, cut worms mole crickets
- Check Nursery Cooperative Website for up to date labels and 24 C labels.

Nursery Insects



"I wouldn't do that, bartender. ... Unless, of course, you think you're fast enough."